



**SCIENTIFIC COMMITTEE  
FIFTH REGULAR SESSION**

10-21 August 2009  
Port Vila, Vanuatu

---

**Mitigation Measures For Pelagic Longline Gear: A Report On The Work Of The Seabird  
Bycatch Working Group, Agreement On The Conservation Of Albatrosses And Petrels**

---

**WCPFC-SC5-2009/EB- IP-9**

**ACAP<sup>1</sup>**

---

<sup>1</sup> Agreement on the Conservation of Albatrosses and Petrels (ACAP)



# **MITIGATION MEASURES FOR PELAGIC LONGLINE GEAR: A REPORT ON THE WORK OF THE SEABIRD BYCATCH WORKING GROUP, AGREEMENT ON THE CONSERVATION OF ALBATROSSES AND PETRELS**

**Author: Agreement on the Conservation of Albatrosses and Petrels (ACAP)**

## **Abstract**

ACAP's Seabird Bycatch Working Group recently assessed the suitability of pelagic mitigation technologies for future research, and reviewed seabird bycatch mitigation measures for pelagic longline fishing to identify knowledge gaps. The products of this work are two tables (Tables 1 and 2), which have been endorsed by ACAP as representing the current best scientific advice. The literature review of mitigation measures showed that some of the measures currently listed by some RFMOs would benefit from further development and testing as they currently have little empirical support on their efficacy.

## **Introduction**

The Agreement on the Conservation of Albatrosses and Petrels (ACAP) is an international Agreement that aims to achieve and maintain a favourable conservation status for albatrosses and petrels, perhaps the most threatened group of birds in the world. While these seabirds face threats both on land and at sea, the greatest threat to their survival is widely acknowledged to be incidental mortality in commercial fisheries, particularly those using longline and trawl gear types.

In recognition of the serious problem posed to seabirds by fisheries interactions, ACAP's Advisory Committee has established a Seabird Bycatch Working Group (SBWG). This working group had been formed to advise the Agreement on actions that will assist in assessment, mitigation and reduction of negative interactions between fishing operations and albatrosses and petrels. The working group comprises representatives from ACAP's 13 Parties, together with invited experts with relevant technical or other expertise. The SBWG has met twice since 2007, and copies of the reports of its meetings can be found at <http://www.acap.aq>

This paper provides a summary of issues relating to bycatch mitigation that may be of use to the WCPFC in developing research and management approaches to mitigate seabird bycatch in its fisheries.

## **Review of Pelagic Longline Mitigation Measures**

The SBWG considers that interactions with pelagic fisheries managed by RFMOs arguably constitute the largest conservation threat to seabirds in the southern oceans. Although several seabird avoidance measures have been trialled to varying degrees in pelagic fisheries, proven and accepted seabird avoidance measures require substantial improvement.

In order to progress the development of relevant mitigation research, the Working Group has developed a plan of research for pelagic longline fisheries, including identifying specific research experiments needed, principal investigators, best host locations, and possible funding sources. It approached this task in two ways, in the light of new data provided to the working group and further expert opinion.

1. An assessment of the suitability of pelagic mitigation technologies for future research and application was carried out. Mitigation measures were grouped as primary, secondary, or other, and a priority ranking for future research assigned on a 5 point scale. Primary measures were those considered likely to be effective without other mitigation measures, and secondary measures were those considered useful for deployment with other measures, but unlikely to significantly reduce bycatch if used in isolation. The results of this assessment are shown in Table 1, together with details of the criteria used for assessment.
2. Seabird bycatch mitigation measures for pelagic longline fishing were reviewed and knowledge gaps identified. The review was based on published literature and expert opinion. The results of the review are shown in Table 2.

Tables 1 and 2 have been endorsed by ACAP as representing the current best scientific advice. The WCPFC and its Members are encouraged to use these materials to guide the development of policy and practice within fisheries under their jurisdiction.

### **Priorities for Research (Table 1)**

It was assessed that from a global research perspective, bird scaring lines, the bait setting capsule and side setting were the highest priority for research. Weighted branchlines, the bait pod, smart hooks and circle hooks were high priorities; and blue dyed squid was of moderate priority. Research on technologies such as the underwater setting chute, night setting, line shooters, thawed bait, strategic offal discharge, blue-dyed fish, fish oil and bait casting machines, were considered a lower priority and were not discussed further. With respect to night setting, the Working Group acknowledged the effectiveness of this mitigation measure, but believed further research on this was not needed.

The Working Group agreed that seabird bycatch mitigation research should best be carried out in locations where seabird interactions with pelagic gear are most intense, as it is these locations that would yield the most useful research outcomes. Locations where aggressive species are most abundant and overlap with fisheries include the pelagic fisheries of Chile in winter, Uruguay and Brazil from May through September, and in South Africa in winter. Personnel from BirdLife International's Albatross Task Force are currently in place in Chile, Brazil, Uruguay, South Africa and Namibia where they are currently collaborating with fishers in seabird bycatch mitigation research programs.

Specific research projects were identified that may be of relevance for WCPFC pelagic longline fisheries. Australia has led the development of the bait setting capsule, a device designed to deliver baited hooks to a depth beyond the access of foraging seabirds at the stern of a pelagic longline vessel. Graham Robertson (Australian Antarctic Division) has funding to develop a prototype and carry out pilot research to demonstrate the efficient performance of a prototype underwater setting capsule. Pending a positive outcome of pilot research, Dr. Robertson is seeking funding to carry out comprehensive research to determine the relative performance of the bait setting capsule, side setting and conventional stern setting. A location to stage this research effort has not been established at this stage.

Ed Melvin ( Washington Sea Grant) is developing a streamer line system for pelagic longline fisheries and has plans to trial the streamer line system in two "worst case" southern hemisphere, pelagic fisheries. Funding is in place to carry out this research. Trials will compare the relative efficiency of the streamer line designed to a control of no deterrent and to a second mitigation technology to be determined. The host locations will include South Africa and either Brazil, Chile or Uruguay. Work is scheduled to be completed in 2009.

Fisheries Managers in New Zealand and Australia have procured “safe leads”, a product which promises to eliminate safety issues related to weighted branchlines. Pilot-level testing of these weights within Australian and New Zealand fisheries has already been undertaken.

### **Seabird bycatch mitigation measures for pelagic longline fishing (Table 2)**

The Seabird Bycatch Working Group recommended that its advice on current best practice mitigation, including the application of combinations of measures (Table 2) be provided to the WCPFC and other relevant RFMOs.

It should be noted that many of the mitigation measures currently adopted by fishers and fisheries managers have little empirical support as to their efficacy. This applies to measures such as side setting, light tori lines, bait casting machines, blue-dyed bait and line-shooter effect on mainline tension.

At SBWG— 2 there was considerable discussion on the use of light tori or bird scaring lines, a variation on the conventional tori line. This measure has recently been proposed as an effective mitigation measure for pelagic longline fisheries. The Working Group noted that there was conflicting information on the effectiveness of this measure. Light tori lines (short streamers and no drag) have been used by the foreign Asian fleet operating in South African waters where substantial seabird bycatch has been reported (0.44 birds killed per 1000 hooks). As a result of improved compliance in 2008, these vessels began using conventional tori lines and seabird bycatch was reduced to 0.05 birds per 1000 hooks. This is likely to be due to a number of factors, but anecdotal evidence suggests that the improved tori line design is a substantial contributor.

A recent study by Yokota et al (2008) tested conventional and light bird scaring lines and compared the frequency of bait-taking behaviour by Laysan albatrosses for each type of bird scaring line. A similar study conducted by Brouwen et al (2008) contained confounding effects and inadequate description of methodologies. Hence it is not possible to draw confident conclusions from this study. Other information from Neves et al (2008) indicates that light bird scaring lines significantly reduced seabird mortality in the absence of any other mitigation measures.

SBWG members found the evidence for effectiveness in the Yokota study to be unconvincing because of the small number of sets (18) in one experiment and the fact that no albatrosses were caught when either bird scaring line type was in use. In a second experiment, although a significant difference in seabird mortality between the two types of bird scaring lines was detected, the confidence limits around the mean values of both treatments overlapped extensively. The SBWG concluded that thorough comparative experimental assessment of light and conventional bird scaring lines needs to be undertaken against Southern Ocean assemblages of diving seabirds (e.g., *Procellaria* sp petrels and *Puffinus* sp. shearwaters) and albatrosses, with research based on larger sample sizes and more transparent methodologies before the measure could be applied with any confidence.

### **Information on ACAP species that occur within the WCPFC Convention Area**

Over the last 18 months a series of assessments has been produced for all of the species on Annex 1 of the Agreement. These provide comprehensive data on the population status, trends and distribution of albatrosses and petrels, including species that occur within the WCPFC convention area such as the black-browed, black-footed, grey-headed, short-tailed, Laysan and wandering albatrosses. These species assessments available on the ACAP website and can be downloaded from <http://www.acap.aq>

## **Recommendations**

It is recommended that the WCPFC:

1. Reviews the information provided in Table 1 when considering the application of currently available mitigation methods;
2. Strongly encourages Members to collaborate on implementing the research initiatives outlined in Table 1.

**Table 1. Assessment of the suitability of pelagic mitigation technologies for future research and application. Rankings have been assigned on a 5 point scale, where 5 is the highest ranking. See below for details of the criteria used for assessment.**

<b>Mitigation</b>	Effective surface feeding birds	Effective diving birds	Practical	Safe	Cost Capital	Cost Ops	DWF/ Dom	Compliance	Future Research Priority
<b>Primary</b>									
Streamer lines	4	3	4	4	5	5	5/5	1	5
Weighted branchlines	4	3	5	1	4	4	5/5	5	4
<b>Underwater Setting</b>									
Chute	2	1	2	3	2	5	1/5	1	1
Bait setting capsule	5	4*	4	4	2	5	5/5	3	5
Bait Pod / Smart hooks	5	4*	3	4*	4	4	5/5	1	4
Night Setting	4	3	5	4	5	3*	5/5	3	1
<b>Secondary</b>									
Circle Hooks	?	?	5	5	5	5	5/5	5	4
Bait placement/casting	2*	2*	5	3	4	4	5/5	1	1
Line shooter?	2	2	5	4	4	4	5/5	1	1
Thawed bait	2	2	3	5	5	5	5/5	1	1
Strategic offal discharge	2	2	3	5	5	5	5/5	1	1
<b>Other</b>									
Side Setting	2*	2*	3	4	4	5	5/5	5	5
Blue Dyed Squid	3	3	3	5	5	4	5/5	1	3
Blue Dyed Fish	1	1	3	5	5	4	5/5	1	1
Fish Oil	1	4	2	4	4	3	5/5	1	2

Each mitigation method was grouped as primary, secondary, or other. Primary measures were those considered likely to be effective without other mitigation measures, and secondary measures were those considered useful for deployment with other measures, but may not significantly reducing bycatch if used in isolation. Side setting, blue-dyed fish and squid bait, and fish oil were regarded as possible candidates for primary mitigation but were considered separately due to their early stage of development and/or limited research results to date. Acoustic alarms, water jets, time-area closures, and artificial lures/bait were not considered. Each was assigned a priority ranking for future research based on the scientific literature and individual experience using the following criteria:

- Effectiveness on surface foraging seabirds
- Effectiveness on diving seabirds
- Practical use on the vessel
- Safe use on the vessel
- Capital Cost – costs for purchase of a specific technology
- Operational Cost – costs related to vessel operations (lost fishing time)
- Applicability to distant water fleets and domestic fleets
- Compliance – the ability to monitor use and performance

Each method was ranked for each criterion on a relative scale of 1 to 5, with 1 being the lowest ranking and 5 being the highest. Considering the ranking for each criterion, each mitigation method was ranked in a similar way resulting in a prioritized list of mitigation methods to focus future research.



**Table 2: Review of Seabird Bycatch Mitigation Measures for Pelagic Longline Fisheries.**

<b>Measure</b>	<b>Scientific evidence for effectiveness in pelagic fisheries</b>	<b>Caveats /Notes</b>	<b>Need for combination</b>	<b>Research needs</b>	<b>Minimum standards</b>
<b>Night setting</b>	Duckworth 1995; Brothers et al. 1999; Gales et al 1998; Klaer & Polacheck 1998; Brothers et al. 1999; McNamara et al. 1999; Gilman et al. 2005; Baker & Wise 2005.	Less effective during full moon, under intensive deck lighting or in high latitude fisheries in summer. Less effective on nocturnal foragers e.g. White-chinned Petrels (Brothers et al. 1999; Cherel et al. 1996).	Recommend combination with bird scaring lines and/or weighted branch lines	Data on current time of sets by WCPFC fisheries. Effect of night sets on target catch for different fisheries.	Night defined as nautical dark to nautical dawn
<b>Side setting</b>	Brothers & Gilman 2006; Yokota & Kiyota 2006.	Only effective if hooks are sufficiently below the surface by the time they reach the stern of the vessel. In Hawaii, side-setting trials were conducted with bird curtain and 45-60g weighted swivels placed within 0.5m of hooks. Japanese research concludes must be used with other measures (Yokota & Kiyota 2006).	Must be combined with other measures. Successful Hawaii trials use bird curtain plus weighted branch lines. In Southern Hemisphere, strongly recommend use with bird scaring lines until side-setting is tested in the region.	Currently untested in the Southern Ocean against seabird assemblages of diving seabirds and albatrosses - urgent need for research. In Japan, NRIFSF will continue testing in 2007.	In Hawaii, side setting is used in conjunction with a bird curtain and 45 weighted swivel within 1m of the baited hook. Clear definition of side setting is required. Hawaiian definition is a minimum of 1 m forward of the stern.

Measure	Scientific evidence for effectiveness in pelagic fisheries	Caveats /Notes	Need for combination	Research needs	Minimum standards
<b>Single bird scaring lines - conventional configuration</b>	Imber 1994; Uozumi & Takeuchi 1998; Brothers et al. 1999; Klaer & Polacheck 1998; McNamara et al. 1999; Boggs 2001; CCAMLR 2002; Minami & Kiyota 2004. Melvin 2003.	Effective only when streamers are positioned over sinking baits. In pelagic fisheries, baited hooks are unlikely to sink beyond the diving depths of diving seabirds within the 150 m zone of the bird scaring line, unless combined with other measures such as line weighting or underwater setting. Entanglement with fishing gear can lead to poor compliance by fishers and design issues need to be addressed. In crosswinds, bird scaring line must be deployed from the windward side to be effective.	Effectiveness increased when combined with other measures e.g. weighted branch lines and/or night setting	Optimal design for pelagic fisheries under development: refine to minimise tangling, optimise aerial extent and positioning, and ease hauling/retrieval. Two studies in progress developing optimal bird scaring line for pelagic fisheries including Washington Sea Grant and Global Guardian Trust in Japan. Controlled studies demonstrating their effectiveness in pelagic fisheries remain very limited.	Current minimum standards for pelagic fisheries are based on CCAMLR Conservation Measure 25-02
<b>Single bird scaring line - Light configuration</b>	Yokota et al. 2008 compared conventional and light bird scaring lines against Laysan albatrosses and considered light lines to be more effective in reducing bait take. A similar study conducted by Brouwer et al. 2008 in New Zealand contained confounding effects	Evidence for effectiveness in Yokota et al (2008) is unconvincing because of small number of sets (18), no seabirds were caught in one experiment, and although a significant difference was detected in a 2 <sup>nd</sup> experiment, the confidence limits around the mean values of both treatments overlapped extensively.		Thorough comparative experimental assessment of light and conventional bird scaring lines against Southern Ocean seabird assemblages of diving seabirds and albatrosses urgently needed. Research needs to be based on larger sample sizes and more transparent methodologies.	Use of this measure is not recommended at this time.

Measure	Scientific evidence for effectiveness in pelagic fisheries	Caveats /Notes	Need for combination	Research needs	Minimum standards
	and inadequate description of methodologies; these concerns preclude confident conclusions to be drawn from this study. Neves et al. 2008 showed light BSLs significantly reduced seabird mortality in the absence of any other mitigation measures.				
<b>Paired bird scaring line – conventional configuration</b>	Two streamer lines best in crosswinds to maximise protection of baited hooks (Melvin et al. 2004).	Potentially increased likelihood of entanglement - see above. Development of a towed device that keeps gear from crossing surface gear essential to improve adoption and compliance.	Effectiveness will be increased when combined with other measures. Recommend use with weighted branch lines and/or night setting	Development and trialling of paired streamer line systems for pelagic fisheries.	Current minimum standards for pelagic fisheries are based on CCAMLR Conservation Measure 25-02
<b>Weighted branch lines</b>	Brothers 1991; Boggs 2001; Sakai et al. 2001; Brothers et al. 2001; Anderson & McArdle 2002; Gilman et al. 2003a; Robertson 2003; Lokkeborg & Robertson 2002, Hu et al. 2005.	Supplementary measure. Weights will shorten but not eliminate the zone behind the vessel in which birds can be caught. Even in demersal fisheries where weights are much heavier, weights must be combined with other mitigation measures (e.g. CCAMLR Conservation Measure 25-02).	Must be combined with other measures e.g. bird scaring lines and/or night setting	Mass and position of weight both affect sink rate. Further research on weighting regimes needed. Testing of safe-leads in progress. Where possible, effect on target catch as well as seabird bycatch should be evaluated. Factors such as swivel weights, mainline tension, bait hooking	Global minimum standards not yet established. Requirements now vary by fishery and vessel. Hawaii minimum requirements are 45g less than 1 m from hook. Australia requires 60 or 90g located 3.5 or 4 m from the hook, respectively, which is a compromise

Measure	Scientific evidence for effectiveness in pelagic fisheries	Caveats /Notes	Need for combination	Research needs	Minimum standards
				position, bait size and life status, deployment position (effect of propeller turbulence) all affect sink rate and need to be quantified.	specification recognising that live bait is used extensively in fishery.
<b>Blue dyed bait</b>	Boggs 2001; Brothers 1991; Gilman et al. 2003a; Minami & Kiyota 2001; Minami & Kiyota 2004; Lydon & Starr 2005. Cocking et al. 2008.	New data suggests only effective with squid bait (Cocking et al. 2008). Onboard dyeing requires labour and is difficult under stormy conditions. Results inconsistent across studies.	Must be combined with bird scaring lines or night setting	Need for tests in Southern Ocean.	Mix to standardized colour placard or specify (e.g. use 'Brilliant Blue' food dye (Colour Index 42090, also known as Food Additive number E133) mixed at 0.5% for minimum of 20 minutes)
<b>Line shooter effect on mainline tension</b>	Reduced bycatch of Northern Fulmar in trials of mitigation measures in North Sea, Lokkeborg & Robertson 2002; Lokkeborg 2003. Increased seabird bycatch in Alaska (Melvin et al. 2001). Robertson et al (2008) found no effect on sink rates in demersal IWL gear.	Supplementary measure. No published data for pelagic fisheries. May enhance hook sink rates in some situations but unlikely to eliminate the zone behind the vessel in which birds can be caught. More data needed. Found ineffective in trials in North Pacific demersal longline fishery (Melvin et al. 2001).	Must be combined with other measures such as night setting and/or bird scaring lines or weighted branch lines	Data needed on effects on hook sink rates in pelagic fisheries.	Not established

Measure	Scientific evidence for effectiveness in pelagic fisheries	Caveats /Notes	Need for combination	Research needs	Minimum standards
	Robertson et al (In Prep) indicates that use of a line shooter in pelagic longline fisheries to reduce mainline tension (e.g., for deep setting) slows significantly the sink rates of hooks.				
<b>Bait caster</b>	Duckworth 1995; Klaer & Polacheck 1998.	Not a mitigation measure unless casting machines are available with the capability to control the distance at which baits are cast. This is necessary to allow accurate delivery of baits under a bird scaring line. Needs more development. Few commercially-available machines have this capability.	Not recommended as a mitigation measure.		
<b>Underwater setting chute</b>	Brothers 1991; Boggs 2001; Gilman et al. 2003a; Gilman et al. 2003b; Sakai et al. 2004; Lawrence et al. 2006.	For pelagic fisheries, existing equipment not yet sturdy enough for large vessels in rough seas. Problems with malfunctions and performance inconsistent (e.g. Gilman et al. 2003a and Australian trials cited in Baker & Wise 2005)	Not recommended for general application	Design problems to overcome	Not yet established

<b>Measure</b>	<b>Scientific evidence for effectiveness in pelagic fisheries</b>	<b>Caveats /Notes</b>	<b>Need for combination</b>	<b>Research needs</b>	<b>Minimum standards</b>
<b>Management of offal discharge</b>	McNamara et al. 1999; Cherel et al. 1996.	Supplementary measure. Definition essential. Offal attracts birds to vessels and where practical should be eliminated or restricted to discharge when not setting or hauling. Strategic discharge during line setting can increase interactions and should be discouraged. Offal retention and/or incineration may be impractical on small vessels.	Must be combined with other measures.	Further information needed on opportunities and constraints in pelagic fisheries (long and short term).	Not yet established for pelagic fisheries. In CCAMLR demersal fisheries, discharge of offal is prohibited during line setting. During line hauling, storage of waste is encouraged, and if discharged must be discharged on the opposite side of the vessel to the hauling bay.
<b>Thawing bait</b>	Brothers 1991; Duckworth 1995; Klaer & Polacheck; Brothers et al 1999.	Supplementary measure. If lines are set early morning, full thawing of all bait may create practical difficulties.	Must be combined with other measures.	Evaluate sink rate of partially thawed bait.	

## REFERENCES

- Anderson, S. and McArdle, B., 2002. Sink rate of baited hooks during deployment of a pelagic longline from a New Zealand fishing vessel. *New Zealand Journal of Marine and Freshwater Research* 36, 185–195.
- Baker, G. B., and Wise, B. S. 2005. The impact of pelagic longline fishing on the flesh-footed shearwater *Puffinus carneipes* in Eastern Australia. *Biological Conservation* 126:306 - 316.
- Boggs, C.H., 2001. Deterring albatrosses from contacting baits during swordfish longline sets. In: Melvin, E., Parrish, J.K. (Eds), *Seabird Bycatch: Trends, Roadblocks and Solutions*. University of Alaska Sea Grant, Fairbanks, Alaska, pp. 79–94.
- Brothers, N. and Gilman, E. 2006. Technical assistance for Hawaii-based pelagic longline vessels to modify deck design and fishing practices to side set. Prepared for the National marine Fisheries Service Pacific Islands Regional Office. Blue Ocean Institute, September 2006.
- Brothers, N. P. 1991. Approaches to reducing albatross mortality and associated bait loss in the Japanese long-line fishery. *Biological Conservation*. 55, 255-268.
- Brothers, N., Gales, R. and Reid, T. 1999. The influence of environmental variables and mitigation measures on seabird catch rates in the Japanese tuna longline fishery within the Australian Fishing Zone 1991-1995. *Biological Conservation* 88:85-101.
- Brothers, N., Gales, R., Reid, T., 2001. The effect of line weighting on the sink rate of pelagic tuna longline hooks, and it's potential for minimising seabird mortalities. *CCSBT-ERS/0111/53*.
- Brouwer, S. and Walker, N. 2008. Use of light streamer lines and line weighting on longline vessels and the implications for seabird bycatch. WCPFC Scientific Committee Fourth Regular Session, 11-22 August 2008 WCPFC-SC4-2008/EB-IP-3.
- CCAMLR, 2002. Report of the working group on fish stock assessment. Report of the twenty-first meeting of the Scientific Committee of the Commission for the Conservation of Marine Living Resources. Commission for the Conservation of Marine Living Resources, Hobart.
- Cherel, Y., Weimerskirch, H. and Duhamel., G 1996. Interactions between longline vessels and seabirds in Kerguelen Waters and a method to reduce seabird mortality. *Biological Conservation* 75:63-70.
- Cocking, L.J., Double, M.C., Milburn, P.J. and Brando, V.E. 2008. Seabird bycatch mitigation and blue-dyed bait: A spectral and experimental assessment. *Biological Conservation*, doi:10.1016/j.biocon. 2008.03.003
- Duckworth, K., 1995. Analysis of factors which influence seabird bycatch in the Japanese southern bluefin tuna longline fishery in New Zealand waters, 1989–1993. *New Zealand Fisheries Assessment Research Document* 95/26.
- Gales, R., Brothers, N. and Reid, T. 1998. Seabird mortality in the Japanese tuna longline fishery around Australia, 1988-1995. *Biological Conservation*. 86,37 56.
- Gilman, E., Brothers, N., Kobayashi, D. R., Martin, S., Cook, J., Ray, J., Ching, G., Woods, B. 2003a. Performance assessment of underwater setting chutes, side setting, and blue-dyed bait to minimize seabird mortality in Hawaii longline tuna and swordfish fisheries. Final report. Western Pacific Regional Fishery Management Council. Honolulu, Hawaii, USA. 42pp.

- Gilman, E., C. Boggs, and N. Brothers. 2003b. Performance assessment of an underwater setting chute to mitigate seabird bycatch in the Hawaii pelagic longline tuna fishery. *Ocean and Coastal Management* 46(11-12): 985-1010.
- Gilman, E., N. Brothers, D. Kobayashi. 2005. Principles and approaches to abate seabird bycatch in longline fisheries. *Fish and Fisheries* 6: 35-49.
- Hu, F., Shiga, M., Yokota, K., Shiode, D., Tokai, T., Sakai, H., Arimoto, T. 2005. Effects of specifications of branch line on sinking characteristics of hooks in Japanese tuna longline. *Nippon Suisan Gakkaishi* 71 (1): 33-38.
- Imber, M.J., 1994. Report on a tuna long-lining fishing voyage aboard Southern Venture to observe seabird by-catch problems. Science & Research Series 65. Department of Conservation, Wellington, New Zealand.
- Klaer, N. and T. Polacheck. 1998. The influence of environmental factors and mitigation measures on by-catch rates of seabirds by Japanese longline fishing vessels in the Australian region. *Emu* 98:305-16.
- Lawrence, E., Wise, B., Bromhead, D., Hindmarsh, S., Barry, S., Bensley, N. and Findlay, J. 2006. Analyses of AFMA seabird mitigation trials – 2001 to 2004. Bureau of Rural Sciences. Canberra.
- Lokkeborg, S. and Robertson, G., 2002. Seabird and longline interactions: effects of a bird-scaring streamer line and line shooter on the incidental capture of northern fulmars *Fulmarus glacialis*. *Biological Conservation* 106, 359–364.
- Lokkeborg, S., 2003. Review and evaluation of three mitigation measures - bird-scaring line, underwater setting and line shooter - to reduce seabird bycatch in the north Atlantic longline fishery. *Fisheries Research* 60, 11–16.
- Lydon, G. and Starr, P., 2005. Effect of blue dyed bait on incidental seabird mortalities and fish catch rates on a commercial longliner fishing off East Cape, New Zealand. Unpublished Conservation Services Programme Report, Department of Conservation, New Zealand. 12 p.
- McNamara B, Torre L, Kaaialii G. Hawaii longline seabird mortality mitigation project. Honolulu, HI, USA: Western Pacific Regional Fishery Management Council, 1999.
- Melvin, E. F., B. Sullivan, G. Robertson, and B. Wienecke. 2004. A review of the effectiveness of streamer lines as a seabird bycatch mitigation technique in longline fisheries and CCAMLR streamer line requirements. *CCAMLR Sci.* 11:189-201.
- Melvin, E. F., J. K. Parrish, K. S. Dietrich, and O. S. Hamel. 2001. Solutions to seabird bycatch in Alaska's demersal longline fisheries. Project A/FP-7, WSG-AS 01-01, Washington Sea Grant.
- Melvin, E.F. 2003. Streamer lines to reduce seabird bycatch in longline fisheries. Washington Sea Grant Program, WSG-AS 00-33.
- Minami, H. and Kiyota, M. 2001. Effect of blue-dyed bait on reducing incidental take of seabirds. *CCSBT-ERS/0111/61*. 7pp.
- Minami, H. and Kiyota, M., 2004 . Effect of blue-dyed bait and tori-pole streamer on reduction of incidental take of seabirds in the Japanese southern bluefin tuna longline fisheries. *CCSBT-ERS/0402/08*.



- Neves T.S., Bugoni, L., Monteiro, D.S., Estima, S.C. (2008). Medidas mitigadoras para evitar a captura incidental de aves marinhas em espinhéis no Brasil. Projeto Albatroz and NEMA. In press.
- Robertson, G. 2003. Fast-sinking lines reduce seabird mortality in longline fisheries. Australian Antarctic Division, Tasmania.
- Sakai, H., Fuxiang, H., Arimoto, T., 2004. Underwater setting device for preventing incidental catches of seabirds in tuna longline fishing, CCSBT-ERS/0402/Info06.
- Sakai, H., Hu, F., Arimoto, T. 2001. Basic study on prevention of incidental catch of seabirds in tuna longline. CCSBT-ERS/0111/62.
- Uozumi, Y. and Takeuchi, Y. 1998. Influence of tori pole on incidental catch rate of seabirds by Japanese southern bluefin tuna longline fishery in high seas. CCSBT-WRS/9806/9 revised. 5pp.
- Yokota, K. and Kiyota, M. 2006. Preliminary report of side-setting experiments in a large sized longline vessel. WCPFC-SC2-2006/EB WP-15. Paper submitted to the Second meeting of the WCPFC Ecosystem and Bycatch SWG. Manila, 10th August 2006
- Yokota, K., Minami, H. and Kiyota, M (2008). Direct comparison of seabird avoidance effect between two types of tori-lines in experimental longline operations. WCPFC Scientific Committee Fourth Regular Session, 11-22 August 2008 WCPFC-SC4-2008/EB-WP-7.